

Product specification ANTIBODY

2025-07-18

Anti-h CA242 NA002601 SPTN-5

Product overview

Catalog number C-10-0065

Specificity Antibody recognizes human CA242 (Carbohydrate Antigen 242)

DescriptionMonoclonal mouse antibody cultured in vitro under conditions free from

animal-derived components.

Product buffer solution 50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.1% ProClinTM 300 as a preservative

Shelf life and storage Unspecified, storage at 2–8 °C

Subclass IgG₁

Analyte description CA242 is a kind of sialic acid carbohydrate and one of the glycosphingolipid

tumor markers. CA242 has a relatively high diagnostic sensitivity for pancreatic cancer and rectal cancer. It could also be used for the differential diagnosis of lung cancer, evaluating the prognosis, monitoring the effect of tumor treatment

电话: 021 68119180, 68119181, 68119105

邮箱: medixchina@medixbiochemica.com

网址:www.cnmedixbiochemica.com

and the recurrence of tumors.

Parameters tested on each lot

Product appearance Liquid, may turn slightly opaque during storage

Product concentration 5.0 mg/ml (+/- 10%)

Immunoreactivity 80–120% compared to the reference sample in a CLIA test

IEF Profile Not Determined (N/D)

Purity $\geq 95 \%$

Kinetic parameters

Association rate constant N/D

Dissociation rate constant N/D

Affinity constant N/D

Determination method -

Determination antigen -





Product specification ANTIBODY

2025-07-18

Cross-reactivities

Epitope N/D

Pair recommendations

		DETECTION	
		NA002601	NA002602
CAPT URE	NA002601	-	+
	NA002602	+	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested ELISA, CLIA

C-40-0081 Carbohydrate antigen 242, CA242 **Antigens tested**

Product stability TEMPERATURE, TIME RESULT

-70 °C, 21 days N/D -20 °C, 21 days N/D +4 °C, 21 days N/D +35 °C, 21 days N/D +45 °C, 7 days N/D

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2-8 °C in the product buffer.

Miscellaneous

References



电话: 021 68119180, 68119181, 68119105