

## Anti-h CRP 6402 SPTN-5

### Product overview

---

<b>Catalog number</b>	100145
<b>Specificity</b>	Antibody recognizes human C-reactive protein (CRP), binding is Ca <sup>2+</sup> independent
<b>Description</b>	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	24 months from manufacturing at 2–8 °C
<b>Subclass</b>	IgG <sub>1</sub>
<b>Analyte description</b>	CRP is a member of the class of acute-phase reactants, as its levels rise dramatically during inflammatory processes occurring in the body. CRP rises up to 50,000-fold in acute inflammation, such as infection. It rises above normal limits within 6 hours, and peaks at 48 hours. Its half-life is constant, and therefore its level is mainly determined by the rate of production (and hence the severity of the precipitating cause). CRP is used mainly as a marker of inflammation. Measuring and charting CRP values can prove useful in determining disease progress or the effectiveness of treatments.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/- 10 %)
<b>Immunoreactivity</b>	80–120 % compared to the reference sample in an FIA test
<b>IEF Profile</b>	6.6–7.7
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	$7.3 \times 10^5$ 1/Ms
<b>Dissociation rate constant</b>	$1.2 \times 10^{-4}$ 1/s
<b>Affinity constant</b>	$K_A = 6.2 \times 10^9$ 1/M; $K_D = 1.6 \times 10^{-10}$ M (= 0.16 nM)
<b>Determination method</b>	SPR analysis (ProteOn XPR36)
<b>Determination antigen</b>	Human CRP, Scripps (Cat C0124, Lot 2200701)



#### Legal disclaimer

**Cross-reactivities** Does not recognize serum amyloid-P complex

**Epitope** Not Determined (N/D)

**Pair recommendations**

		DETECTION				
		6402	6403	6404	6405	6407
CAPTURE	6402	+	+	+	+	+
	6403	+	+	+	+	-
	6404	+	+	+	+	+
	6405	+	-	+	+	+
	6407	+	-	+	+	+

Following pairs are especially recommended for the below mentioned assays:

LF: 6402 (membrane) – 6402 (particles), 6402 – 6404, 6402 – 6405

IT: 6405 – 6405, hs-CRP (high sensitivity): 6404 – 6407, 6405 – 6407

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Platforms tested** FIA, LF, IT

**Antigens tested** Recombinant CRP antigen, Lee Biosolutions 140-11R.

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+25 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous** Note that CRP is a pentameric protein. The same antibody can thus be used both as a capture and a detection antibody.

**References** Käpyaho, K., Welin, M.-G., Tanner, P., Kärkkäinen, T. and Weber, T. (1989) Rapid determination of C-reactive protein by enzyme immunoassay using two monoclonal antibodies. Scand. J. Clin. Lab. Invest. 49:389-393



**Legal disclaimer**