

HiDi® DNA Polymerase

Description

HiDi® DNA polymerase is a highly selective DNA polymerase variant, specially evolved for all assays in which **High Discrimination** is required, for instance in allele-specific PCRs, primer extensions or methylation-specific PCRs.

An aptamer based hot-start formulation of the HiDi® DNA polymerase prevents false amplification. Temperatures above 50–55°C cause the aptamer's secondary structure to melt and will set free the polymerase.

HiDi® DNA polymerase efficiently amplifies from primers that are matched at the 3'-end and discriminates primers that are mismatched.

Applications include SNP-detection by allele-specific amplification (ASA) / allele specific PCR, methylation specific PCR (MSP), HLA genotyping and multiplex PCR.

Kit components

Component	S pack*	M pack*
HiDi® DNA Polymerase	50 µL	200 µL
10x HiDi® Reaction Buffer	1 x 1.25 mL	2 x 1.25 mL

*Other pack sizes, bulk orders and customization are available upon request.

Storage and shipment

Transport with cool packs. The reagents should be stored at -20°C upon arrival. The reagents are stable until the expiration date if stored correctly.

Reaction Master Mix set-up

The recommended master mix set-up for a 50 µL reaction volume is shown in the table below.

Reagent	Volume (µL)	Final concentration
HiDi® DNA Polymerase (5 U/µL)	0.5	2.5 U/rxn
HiDi® buffer (10x)	5	1x
∞Forward primer (10 µM)	1	0.2 µM (0.05–1 µM)
∞Reverse primer (10 µM)	1	0.2 µM (0.05–1 µM)
dNTPs (2 mM)	5	200 µM
Template/Sample extract	X	<1000 ng* DNA
Nuclease-free water	Up to 50 µL final volume	

Keep all components on ice.

Spin down and mix all solutions carefully before use.

∞Primers should ideally have a GC content of 40–60% typically.

*Suggested template concentration should be about 10 ng – 1000 ng (genomic DNA) or 1 pg – 1 ng (plasmid/viral DNA) per reaction.

Instrument and program set-up

Cycles	Steps	Temperature	Time
1	Initial denaturation	95°C	2 min
25–40	Denaturation	95°C	15 sec
	Annealing*	54–72°C	30 sec
	Extension	72°C	30 sec /250 bp

*Typically, the annealing temperature is about 3–5°C below the calculated melting temperature of the primers used.



Legal disclaimer

Technical information and support

HiDi® 10x buffer is optimized for short amplicon length (about 60–200 bp). In case of longer amplicons (>500 bp) the addition of magnesium (+ 0.5–1.5 mM) might be needed.

HiDi® DNA polymerase can be used for real-time cycling, by adding a suitable real-time PCR dye.

HiDi® DNA polymerase is a nuclease deficient DNA polymerase, therefore not suitable for hydrolysis probe-based assays. For those assays, HiDi® Taq DNA polymerase (#9201) is recommended.

For technical enquiries or assay development support, please contact us via e-mail at:
mdx@medixbiochemica.com.

Additional information and technical resources are available on our website at:
info.medixbiochemica.com/resources.



Legal disclaimer